



## REVIEW

# Nutrigerontology: why we need a new scientific discipline to develop diets and guidelines to reduce the risk of aging-related diseases

Kris Verburgh

Center Leo Apostel for Interdisciplinary Studies (CLEA), Free University of Brussels (VUB), Brussels, Belgium

## Summary

**Many diets and nutritional advice are circulating, often based on short- or medium-term clinical trials and primary outcomes, like changes in LDL cholesterol or weight. It remains difficult to assess which dietary interventions can be effective in the long term to reduce the risk of aging-related disease and increase the (healthy) lifespan. At the same time, the scientific discipline that studies the aging process has identified some important nutrient-sensing pathways that modulate the aging process, such as the mTOR and the insulin/insulin-like growth factor signaling pathway. A thorough understanding of the aging process can help assessing the efficacy of dietary interventions aimed at reducing the risk of aging-related diseases. To come to these insights, a synthesis of biogerontological, nutritional, and medical knowledge is needed, which can be framed in a new discipline called 'nutrigerontology'.**

**Key words:** aging; aging-related diseases; biogerontology; cardiovascular disease; diabetes type 2; diet; nutrient-sensing pathways; nutrition; obesity; overweight.

## Introduction

Great progress has been made in the course of recent decades in the field of aging research. Biogerontology, the science that studies aging, uncovered some important molecular pathways that are major modulators of the aging process (Kenyon, 2010; Fontana *et al.*, 2010; Bartke *et al.*, 2013). At the same time, society is confronted with an ever-increasing rise in the prevalence of aging-related diseases, such as cardiovascular disease, type 2 diabetes, and Alzheimer's disease. Extending knowledge from biogerontology to clinical medicine and nutritional science can help discover adequate dietary recommendations to prevent or slow down the progress of various aging-related diseases. This knowledge can be framed into 'nutrigerontology', a new scientific discipline that encompasses biogerontology, medicine, and dietetics. Nutrigerontology researches how food substances, foods, food

patterns, and diets influence the risk of aging-related diseases and (healthy) lifespan. Nutrigerontology can also be considered as a practical application of biogerontology. This article wants to clarify how biogerontological insights can be used to create dietary recommendations that can reduce the risk of various aging-related diseases that are on the rise in both developed and developing countries.

## Reduced insulin/IGF-1 and mTOR signaling increases lifespan

In the field of biogerontology, the two most well-known pathways implicated in the aging process are the insulin/insulin-like growth factor signaling pathway (IIS pathway) (Bartke *et al.*, 2013) and the mammalian or mechanistic target of rapamycin (mTOR) pathway (Johnson *et al.*, 2013). We will initially focus on these two canonical pathways to show how a better understanding of the aging process can help to develop better long-term diet recommendations. The mTOR and IIS pathways are nutrient-sensing pathways, implying that they are activated by nutrients that we eat, such as carbohydrates (which mainly activate the IIS pathway, but also the mTOR pathway) (Bartke *et al.*, 2013) and amino acids (which mainly activate the mTOR pathway, but also the IIS pathway) (Wullschleger *et al.*, 2006). The IIS pathway exerts its effects through transmembrane insulin and IGF-1 receptors, which initiate glucose uptake in the cell, and stimulate cell growth and cell proliferation. The mTOR protein is intracellularly located and a potent stimulator of protein translation.

A mounting body of research shows that loss of function of the IIS pathway and mTOR pathway increases lifespan. Modulating these pathways, genetically or nutritionally, can impact the rate of aging and postpones the advent of aging-related diseases.

For example, mice with a fat-specific insulin receptor knockout (FIRKO) genotype live 18% longer (Blüher *et al.*, 2003) and reducing insulin receptor signaling in the mouse brain extended lifespan up to 18% (Taguchi *et al.*, 2007). Homozygous deletion of the insulin receptor substrate 1 (*Irs1*<sup>-/-</sup>), an effector of the insulin receptor, increased median lifespan by 32% in female mice. However, deletion of *IRS-1* in male mice did not lead to an increased lifespan (Selman *et al.*, 2008). Mice heterozygous for the IGF-1 receptor (*Igf1r*<sup>+/-</sup>) live on average 26% longer than their wild-type littermates (Holzenberger *et al.*, 2003). In humans, the insulin/insulin-like growth factor signaling pathway also is involved in lifespan regulation. Polymorphic variants of IGF-related pathways confer health and lifespan benefits in humans (Bonafé *et al.*, 2003; Kojima *et al.*, 2004; Pawlikowska *et al.*, 2009). Female Ashkenazi Jewish centenarians show an overrepresentation of loss-of-function mutations in the IGF-1 receptor (Suh *et al.*, 2008). Laron dwarves, who are growth hormone receptor deficient, have reduced levels of IGF-1, and despite being generally overweight, they have a major reduction in the risk of type 2 diabetes and cancer (Guevara-Aguirre *et al.*, 2011; Steuerma *et al.*, 2011). People with acromegaly (due to increased growth hormone and IGF-1 production) have a two-

## Correspondence

Kris Verburgh, Center Leo Apostel for Interdisciplinary Studies, Free University of Brussels (VUB), Krijgskundestraat 33, B-1160 Brussels, Belgium. Tel.: +32-2-640.67.37; fax: +32-2-644.07.44; e-mail: kris.verburgh@vub.ac.be

Accepted for publication 30 September 2014

to threefold increase in mortality, mostly because of vascular disease (Clayton, 2003).

Regarding the mTOR pathway, loss-of-function of mTOR or proteins involved in the mTOR pathway doubles the lifespan in the nematode *Caenorhabditis elegans* (Vellai *et al.*, 2003) and extends lifespan in the fruit fly *Drosophila melanogaster* (Kapahi *et al.*, 2004). Mice that express mTOR at 25% of wild-type mice display a 20% increase in median lifespan (Wu *et al.*, 2013). Rapamycin, a pharmacological mTOR inhibitor, extends lifespan with 9% in male and 14% in female mice (Harrison *et al.*, 2009).

## The IIS and mTOR pathways in aging-related disease

How do the IIS and mTOR pathway contribute to the aging phenotype? A continuous stimulation of the IIS and mTOR pathways leads to a higher risk of aging-related diseases and reduced lifespan via reduced autophagy (less clearance of protein aggregates and cellular organelles), increased protein agglomeration and proteotoxicity (mTOR activation leads to increased protein production and reduced protein clearance), inflammation, reduced expression of antioxidant proteins, mitochondrial dysfunction, and other mechanisms (Kenyon, 2010; Fontana *et al.*, 2010; Johnson *et al.*, 2013; Cuervo, 2008). These mechanisms increase the risk of insulin resistance (Shah *et al.*, 2004), atherosclerosis (Martinet *et al.*, 2014), cardiomyopathy (Willis & Patterson, 2013), neurodegenerative diseases (O'Neill *et al.*, 2012, 2013), cancer (Martini *et al.*, 2014), osteoporosis (Glantschnig *et al.*, 2003) and other aging-related diseases and symptoms.

One of the most intriguing aspects of the IIS and mTOR pathways is that these are *nutrient-sensing* pathways, implying that they are activated by nutrients (carbohydrates and amino acids), which are contained in the food we eat. An overabundance of these nutrients can accelerate the aging process and increase the risk of aging-related diseases such as cardiovascular disease and type 2 diabetes. This fits in a new emerging paradigm shift, in which the aging process is viewed as a consequence of continuous cellular stimulation and hyperfunction in adulthood, in which a continuous bombardment of our cells with nutrients, growth factors, and mitogenic stimuli accelerates the aging phenotype (Gems & de la Guardia, 2012; Blagosklonny, 2006).

Extrapolating knowledge from the biogerontological field to nutritional science and medicine could lead to a new scientific field, called 'nutrigenontology'. A diet that reduces the risk of aging-related diseases could be one that delivers carbohydrates, amino acids, and fats in such a way that pro-aging nutrient-sensing pathways such as the IIS and mTOR pathway are less activated. The ideal composition of such a diet remains to be determined, but animal and human studies already provide interesting clues.

## mTOR, diets, and nutrition

Because aging-related pathways like mTOR are activated by amino acids, a diet that does not provide an overabundance of amino acids can be considered beneficial (Efeyan *et al.*, 2012). Restricting proteins is one of the few dietary interventions that can increase lifespan in rodents (Yu *et al.*, 1985; Leto *et al.*, 1976; Ross, 1961; Ross & Bras, 1975). Mice on low-protein diets (5% to 15% protein) live up to 150 weeks compared to 100 weeks for mice on a high-protein diet (50% protein), and this increase in lifespan was associated with reduced mTOR activation (Solon-Biet *et al.*, 2014). Reducing the amount of specific amino acids in the diets extends lifespan in various species. The lifespan of *Drosophila* is extended by caloric restriction, but this effect is nullified when essential

amino acids are provided (Grandison *et al.*, 2009). In yeast cells, the amino acids threonine and valine activate Tor1 (the yeast orthologue of mTOR), promoting aging (Mirisola *et al.*, 2014). Restriction of the essential amino acid methionine increases lifespan in rodents (Richie *et al.*, 1994; López-Torres & Barja, 2008). Vascular aging is accelerated in rats fed a high methionine (2%) or high-protein (50%) diet (Fau *et al.*, 1988). Alzheimer mice that undergo protein restriction cycles (a diet lacking essential amino acids is provided every other week), reduce IGF-1 levels, slow down the deposition of tau protein in the brain, and score higher on behavioral performance tests (Parrella *et al.*, 2013).

How can we use these insights in the context of human diets? Meat is an important source of protein in the human diet in many cultures. Recent large studies show an association with meat intake and cardiovascular disease, type 2 diabetes, cancer, and all-cause mortality (Pan *et al.*, 2012; Rohrmann *et al.*, 2013; Ericson *et al.*, 2013; Gonzalez & Riboli, 2010). However, the increased mortality due to meat intake cannot be solely attributed to the protein content of meat. Meat, and especially red processed meat, contains other ingredients that may have negative effects, such as specific fats, salt, or preservatives (Pan *et al.*, 2012). Even so, a mounting body of research suggests a causal relationship between amino acid intake and a less optimal metabolism in humans. For example, a twofold rise in amino acid blood level leads to a 25% reduction in insulin sensitivity (Krebs *et al.*, 2002) and directly infusing amino acids into the bloodstream increases insulin resistance in humans (Tremblay *et al.*, 2007, 2005). Additionally, recent studies show an association between increased protein intake from meat and aging-related diseases in humans (Ericson *et al.*, 2013; Levine *et al.*, 2014). According to a study conducted in 6318 men and women from the NHANES III survey, people aged 50–65 years with a high protein intake (>20% of total calories) were four times as likely to die of cancer and had a 75% increase in mortality compared to people with a low protein intake (<10% of total calories). Participants with a low protein intake had also significantly lower IGF-1 levels. However, in this study, people over 65 may benefit from an increased protein intake (but only when their IGF-1 was also low). Yet, across all ages, there remained a fivefold increase in diabetes risk when comparing low with high protein intake. It is however important to point out that this study has its limitations: the sample size is rather modest, the food intake data are based on personal food questionnaires which are prone to underreporting, and the researchers do not substantially control for the overall balance of the diet, as people who eat much animal protein tend to consume less fruit or vegetables for example.

Interestingly, in this study, there was no association between *vegetable* protein intake and mortality (Levine *et al.*, 2014). Other studies indeed corroborate that there is a difference between animal and vegetable protein in terms of health. Rats who are fed soy proteins instead of casein proteins have an increased lifespan (Iwasaki *et al.*, 1988). In humans, proteins from vegetable sources induce IGF-1 less than proteins from animal sources (Allen *et al.*, 2002). These physiological differences between animal and vegetable proteins can be explained by the fact that vegetable protein contains less methionine and sulfur-rich amino acids than animal protein. Substituting red meat with healthier (vegetable) proteins sources such as beans, lentils, tofu, and nuts could be recommended (Pan *et al.*, 2012).

Some studies suggest that an increased protein intake can be protective in the elderly, reducing the risk of sarcopenia for example (Houston *et al.*, 2008; Meng *et al.*, 2009). While there is suggestive evidence that a higher protein intake can protect against sarcopenia in the elderly, there is no conclusive evidence that an increased protein intake reduces frailty or lowers overall mortality in older people

(Pedersen & Cederholm, 2014). Interestingly, a low protein intake in the elderly leading to reduced muscle mass is accompanied by a reduction in IGF-1 (Castaneda *et al.*, 2000). Animal and human studies show that reduced IGF-1 signaling leads to lower muscle mass, lower body weight, and a smaller phenotype but an increased lifespan, suggesting that decreased muscle mass is not always detrimental (Bartke *et al.*, 2013; Holzenberger *et al.*, 2003; Bonafè *et al.*, 2003; Kojima *et al.*, 2002; Pawlikowska *et al.*, 2009; Suh *et al.*, 2008; Guevara-Aguirre *et al.*, 2011; Steuerman *et al.*, 2011). Additionally, even if an increased protein intake should be beneficial in the elderly, recent studies suggest that a high protein intake in the nonelderly (<65 years old) can be detrimental, especially with regard to IGF-inducing animal proteins and especially in the long term (>15 years) (Levine *et al.*, 2014). Many studies in humans that assess protein intake and health are short- to medium-term studies.

While amino acids activate mTOR, there exists various substances that inhibit mTOR. Examples of such substances can be found in specific foods considered to confer health benefits, such as the ethylxanthine caffeine in coffee (Reinke *et al.*, 2006; Wanke *et al.*, 2008) and the polyphenol EGCG present in green tea (Van Aller *et al.*, 2011), which reduce mTOR signaling. Quercetin, a substance found in fruits, vegetables, and spices, especially capers, lovage, and apples, is a mTOR inhibitor (Bruning, 2013). Resveratrol, a polyphenol found in the skin of red grapes also inhibits mTOR (Liu *et al.*, 2010). It is important to note that many nutrients that confer potential health benefits do not solely affect mTOR, but often act on various aging-related pathways (see further on), including mTOR.

Other substances in food act downstream of the mTOR receptor. One of the consequences of too much mTOR activation is a reduction in autophagy, which leads to less degradation of protein aggregates in the cell. Protein aggregation is implicated in all kind of aging-related diseases, such as cardiomyopathy and Alzheimer's disease (Choi *et al.*, 2013). Certain nutritional compounds, however, have the potential to attenuate protein aggregation, directly or indirectly. Curcumin, a substance found in the yellow spice turmeric, can directly inhibit the formation of amyloid-beta oligomers by interacting with the amyloidogenic regions of the amyloid-beta protein (Yang *et al.*, 2005; Ono *et al.*, 2004). Oleocanthal, a phenolic substance responsible for the bitter taste in extra-virgin olive oil, can reduce tau fibrillization (Monti *et al.*, 2011) and enhance B-amyloid clearance *in vitro* and *in vivo*, potentially reducing the risk of Alzheimer's disease (Abuznait *et al.*, 2013). Cinnamon extract can reduce B-amyloid formation and slow down cognitive deterioration in Alzheimer's disease model mice (Frydman-Marom *et al.*, 2011).

## The IIS pathway, diets, and nutrition

Besides amino acids, nutrients like carbohydrates stimulate the IIS and mTOR pathway, especially via the insulin receptor, which activates both the mTOR and the IIS pathway (Um *et al.*, 2006; Jewell & Guan, 2013). The glycemic index (GI) measures the area under the blood glucose curve after ingestion of a specific carbohydrate-containing food and indicates how much this food increases blood sugar and subsequently insulin production (Brouns *et al.*, 2005). The glycemic load (GL) is a more precise measurement than the GI, because the GL takes into account both the GI and how much absorbable carbohydrates are contained in a specific food (Foster-Powell *et al.*, 2002). Especially, bakery foods that contain an ample amount of carbohydrates combined with fats increase insulin production (Holt *et al.*, 1997).

The IIS and mTOR pathway are two mechanisms through which high-glycemic index and glycemic load diets exert deleterious effects in the

long term, because such diets deliver high amounts of nutrients in the form of easy-digestible carbohydrates, constituting a nutrient overload that activates aging-related nutrient-sensing pathways and leads to resistance of the insulin receptors (Um *et al.*, 2006). Studies show that diets with a high-glycemic index and load increase the risk of various aging-related diseases such as type 2 diabetes, stroke, and heart disease (Beulens *et al.*, 2007; Salmerón *et al.*, 1997; Liu *et al.*, 2000; Villegas *et al.*, 2007; Sieri *et al.*, 2013). On the other hand, some studies show no effect of glycemic index and load on type 2 diabetes (Mosdøl *et al.*, 2007; Meyer *et al.*, 2000).

Regarding weight loss, low-glycemic index diets are superior compared to low fat diets in terms of improving cardiovascular parameters and weight loss according to a Cochrane meta-analysis (Thomas *et al.*, 2007). Adhering to a low-glycemic index diet improves metabolic parameters more than an isocaloric low-fat diet (Ebbeling *et al.*, 2012). A strict hypocaloric low-glycemic index-load and mainly vegetable-based diet could even reverse diabetes in all subjects in 8 weeks time (Lim *et al.*, 2011).

It should be noted that a large intake of fast-acting carbohydrates can have deleterious effects other than activation of the IIS pathway and mTOR pathway, for example via formation of advanced glycation end products (AGEs) (Singh *et al.*, 2001). However, glycation and cross-linking are also well-known aging-mechanisms in the field of biogerontology (Vitek *et al.*, 1994; Kilhovd *et al.*, 1999).

Not only the macronutrient content (carbohydrates, proteins, or fats) of a diet, but also the micronutrient content can influence metabolism and more specifically insulin sensitivity. For example, blueberries contain anthocyanidins of which the intake is associated with improved insulin sensitivity and a decreased risk of type 2 diabetes (Muraki *et al.*, 2013). Extracts from cinnamon can improve fasting blood glucose levels and lower glycated HbA1c concentrations in type 2 diabetes patients (Lu *et al.*, 2012). Intake of cacao (the main ingredient of dark chocolate) is associated with increased insulin sensitivity and a reduced risk of cardiovascular disease (Buitrago-Lopez *et al.*, 2011) and can slow down the progress of mild cognitive impairment in elderly people (Desideri *et al.*, 2012).

## The role of fat and other nutrient-sensing pathways in aging

In this article, the role of protein and carbohydrates as macronutrients has been discussed. Fat constitutes a third important macronutrient. While amino acids and carbohydrates can directly activate aging-pathways such as mTOR and insulin/IGF-1, the role of specific proteins targeted by fatty acids and their potential role in aging is still unclear (Hou & Taubert, 2012; Hansen *et al.*, 2013). However, mTOR and insulin/IGF-1 signaling play an important role in lipid metabolism during aging. Overstimulation of the mTOR and IIS pathway in preadipocytes by nutrients like carbohydrates and amino acids may contribute to their conversion into senescent-like preadipocytes which secrete pro-inflammatory compounds and induce insulin resistance in surrounding adipose cells. Additionally, senescent-like preadipocytes and insulin-resistant adipocytes are less efficient in handling and processing fatty acids, causing ectopic fatty acid deposition (in the liver, bone marrow, or blood vessels) contributing to heart disease, fatty liver disease, and stroke (Tchkonina *et al.*, 2010; Stout *et al.*, 2014).

As already stated in this paper, mTOR and insulin/IGF-1 signaling are prime examples of nutrient-sensing pathways involved in the aging process. Besides these canonical pathways, several other nutrient and energy sensing pathways are involved in the aging process. An example

is the AMPK pathway, which is activated when the energy status in the cell is low (reduced ATP and increased AMP levels) (Houtkooper *et al.*, 2010). Activated AMPK improves glucose uptake, mitochondrial biogenesis, and  $\beta$ -oxidation of fatty acids (Winder & Hardie, 1999). Other nutrient-sensitive modulators of the aging process are sirtuins (NAD-dependent histone deacetylases), which are activated by a low energy status of the cell (increased NAD<sup>+</sup>) (Houtkooper *et al.*, 2012). Various nutrient-sensing pathways interact with each other. Mice overexpressing the sirtuin SIRT6 live longer and have reduced insulin and IGF-1 signaling (Kanfi *et al.*, 2012), while AMPK signaling reduces mTOR (Alers *et al.*, 2012) and works together with sirtuins to regulate energy metabolism (Price *et al.*, 2012).

## Hormesis

Another interesting biogerontological insight is the concept of hormesis. Hormesis describes how small amounts of toxic substances can be beneficial because they induce cellular stress resistance mechanisms, such as upregulation of heat shock proteins, antioxidant enzymes, or phase II detoxification enzymes (Calabrese *et al.*, 2007). For example, coffee contains slightly toxic compounds that activate Nrf-2, a transcription factor that upregulates cellular antioxidant proteins via the *antioxidant responsive element* (ARE) transcription factor (Boettler *et al.*, 2011). Many vegetables and fruits contain hormetic substances that could have beneficial effects in the long term. Sulforaphane, a compound in cruciferous vegetables like broccoli, induces phase II detoxification enzymes (Hu *et al.*, 2006), a process that can play a role in protecting brain tissue (Trinh *et al.*, 2008) or reducing the risk of cancer (Hayes *et al.*, 2008; Ambrosone *et al.*, 2004). Hormesis can also explain why most antioxidants do not retard the aging process or increase lifespan (Bjelakovic *et al.*, 2007). Large doses of extracorporeal antioxidants such as vitamin A, vitamin E, or beta-carotene downregulate cellular stress resistance proteins because high levels of antioxidants signal to the cell a reduced need for the production of antioxidant proteins and detoxification enzymes, which are in fact much more effective than extracorporeal antioxidants to provide resistance against cellular stress.

## Caloric restriction and fasting

Besides specific macronutrient ratios and micronutrient intake, less nutrients could also reduce the risk of aging-related diseases. Caloric restriction and fasting drastically reduce IIS and mTOR signaling and increase lifespan in many species, ranging from worms to rhesus monkeys (Fontana *et al.*, 2010; Colman *et al.*, 2009). A long-term study (>20 years) at the Wisconsin National Primate Research Center showed that age-related mortality was threefold lower in the caloric-restricted rhesus monkeys (Colman *et al.*, 2009). However, another long-term study conducted by the National Institute of Aging (NIA) in caloric-restricted rhesus monkeys did not show an increase in lifespan (Mattison *et al.*, 2012). This different outcome could be because in the NIA study, the control group food rations had a different dietary composition or because the control group was not fed *ad libitum* and as a consequence was also subjected to a minor form of caloric restriction (Partridge, 2012). This latter explanation is supported by the fact that the control monkeys in the NIA study weighed substantially less than the national average for body weight in the same age groups (Colman *et al.*, 2014). However, the question remains if *ad libitum* feeding is more an obesity model than a normal feeding state in primates, implying that caloric restriction would 'normalize' extra food intake in primates rather than

further improving health and lifespan in primates with a normal food intake. Additionally, other factors besides diet such as the genetic background of the test subjects or the experimental design may also account for these discrepancies.

In humans, long-term caloric restriction is associated with improved cardiometabolic parameters and a reduction of atherosclerosis and diastolic dysfunction (Fontana *et al.*, 2004; Meyer *et al.*, 2006; Heilbronn *et al.*, 2006). Intermittent fasting (also referred to as 'low level caloric restriction') could be a more achievable alternative to caloric restriction. Fasting every other day or on several days per week (during which one consumes up to 75% less calories) leads to a reduced total caloric intake in the long term. Preliminary studies show improvement of various cardiovascular markers such as LDL cholesterol and insulin sensitivity (Varady *et al.*, 2009; Halberg *et al.*, 2005). Fasting could be especially beneficial in the treatment of cancer, as fasting can make tumor cells more vulnerable to chemotherapy and improve regeneration of the immune system after chemotherapy (Raffaghello *et al.*, 2008; Lee *et al.*, 2012; Cheng *et al.*, 2014). Further research is needed to investigate the long-term effects of fasting and caloric restriction on aging-related diseases in humans.

## Discussion

Insights from biogerontology can be very useful to assess the efficacy of diets that aim to reduce the risk of aging-related diseases, or mitigate them. Without biogerontology as a benchmark or touchstone, one has to rely mainly on clinical trials which focus on short- or medium-term primary outcomes, like reducing LDL cholesterol or triglycerides, but do not take into account long-term outcomes, like an increased (healthy) lifespan. A thorough understanding of the aging process can help to better predict the efficacy of diets and specific nutritional interventions. For example, biogerontological knowledge predicts that diets being high in animal protein (like the Atkins diet or some interpretations of the paleo diet) are probably not healthy in the long term and that taking antioxidant dietary supplements does not increase lifespan. On the other hand, specific diets that have been shown to reduce the risk of various aging-related diseases seem to act via important biogerontological pathways and mechanisms.

For example, a Mediterranean diet can substantially reduce the risk of cardiovascular disease, and this to a greater extent than a standard low-fat diet (De Lorgeril *et al.*, 1994). The Mediterranean diet is a low-glycemic index diet that contains many mithormetic, anti-amyloidogenic, and anti-inflammatory compounds found in vegetables, fruits, red wine, olive oil, or nuts. In Okinawa, where substantially more centenarians live with often good health, there exists a tradition of adhering to a vegetable-rich low-glycemic index diet and practicing a form of minor caloric restriction (Okinawans tend to consume small portions) (Willcox *et al.*, 2006, 2007). Vegetarian diets are associated with a reduced all-cause mortality (Orlich *et al.*, 2013). Compared to meat-containing diets, a vegetarian diet provides more plant-based proteins and a lower methionine content, which could contribute to the increase in lifespan and health (McCarty *et al.*, 2009; Cavuoto & Fenech, 2012).

In the future, even healthier diets could be devised using biogerontological insights. These diets would not overstimulate mTOR, insulin, and IGF-1 receptors, would deliver mithormetic substances that stimulate cellular stress resistance pathways and would provide a copious supply of substances that beneficially influence myriad mechanisms involved in the aging process such as protein/amyloid aggregation, autophagy, lipid peroxidation, and mitochondrial dysfunction. A synthesis of biogerontological, nutritional, and medical knowledge can be

framed in a new field called 'nutrgerontology'. Nutrgerontology can be defined as the scientific discipline that studies the impact of nutrients, foods, macronutrient ratios, and diets on lifespan, the aging process, and aging-related diseases.

The goal of nutrgerontology is to research and use compounds, foods, and diets that can reduce the risk of aging-related diseases and increase the (healthy) lifespan. Some interventions can slow down the progress of these aging-related diseases or even reverse them, like type 2 diabetes or cardiovascular disease (Lim *et al.*, 2011; De Lorgeril *et al.*, 1994). While biogerontology mainly focuses on unraveling the mechanisms of aging and gerontology also focuses on the social and psychological aspects of aging, nutrgerontology signifies a practical approach. Nutrgerontology will be mainly practiced by physicians (cardiologists, endocrinologists, oncologists, neurologists, family physicians) and dieticians. Nutrgerontology can be applied in many different medical fields where aging-related diseases are common, such as cardiology (atherosclerosis, diastolic dysfunction, ventricular hypertrophy, high blood pressure), endocrinology (type 2 diabetes), neurology (Alzheimer's disease, stroke, Parkinson's disease), ophthalmology (cataract, aging-related macular degeneration), oncology (cancer), or rheumatology (osteoporosis, sarcopenia, frailty). While geriatrics is the medical discipline that treats the elderly, nutrgerontology is a medical and scientific discipline that wants to act as much as possible on the period before people are old, frail, and afflicted by a myriad of aging-related diseases.

Medical students should be taught about nutrgerontology and learn how various diseases that afflict our society are in fact (accelerated) aging-related diseases. They will learn how evidence-based long-term dietary interventions can reduce the risk of these diseases and mitigate them, slow down their progress or revert them. In a nutrgerontology course, students will be instructed in the aging process and how mechanisms of aging cause many different aging-related diseases. This is important, as addressing and eliminating only one disease of aging, like cardiovascular disease for example, would only add 2.87 years to average lifespan (Barzilai & Rennert, 2012), because patients would still be afflicted by other diseases of aging, like Alzheimer's disease, diabetes, or cancer. Therefore, it would be paramount to address aging itself, and not only specific aging-related diseases without emphasis on aging (biogerontology) and relevant interventions (nutrgerontology).

However, extrapolating knowledge from the biogerontological field to nutritional science and medicine has to be performed carefully, as many model organisms used in aging research (yeast, fruit flies, or mice) are much shorter lived than humans and differ from humans genetically and physiologically and may have been exposed to different mechanisms of natural selection (more selection for fast reproduction in mice for example). On the other hand, the conservation of many aging-related pathways such as mTOR and insulin/IGF-1 in evolutionary very different organisms over millions of years strongly suggests that such pathways are also relevant in humans. Interpretation of study results from model organisms is sometimes challenging, because many confounding factors can influence lifespan in laboratory animals, such as genetic differences between wild-type or laboratory strains, laboratory environment (food supplies, husbandry, bedding, litter mates), and accidental or premature death (Partridge & Gems, 2007).

There is an urgent need of integrating knowledge from the biogerontological field with nutritional science and medicine in view of the increasing epidemic of aging-related and metabolic diseases, driven by a rapidly aging population and an unhealthy lifestyle (Anon, 2011). In 2010, an estimated 524 million people were 65 years or older (8% of

the world population). By 2050, this number will increase to about 1.5 billion (16% of the world population). In 2008, noncommunicable aging-related diseases such as heart disease, Alzheimer's disease, and cancer constituted 86% of the burden of disease in high-income countries. Dementia alone accounted for 600 billion dollar in healthcare costs worldwide. From the age of 65, the risk of dementia doubles every five years resulting in 30% of people between 85–89 years being afflicted with dementia (Anon, 2011).

A better understanding of the aging process and ensuing practical, clinical, and nutritional interventions could not only improve the quality of life and health of many people, but would also reduce aging-related healthcare costs and long-term care expenditures, increase potential working span and productivity, and strengthen economical competitiveness.

Both animal and human research into aging and nutrition should be promoted, so that various ways of how nutrients, foods and diets interfere with the aging process can be researched. Examples of such studies are protein restriction cycles that reduce the progress of Alzheimer's disease in mice or dietary patterns that occur more in centenarian populations (De Lorgeril *et al.*, 1994; Willcox *et al.*, 2007). In light of our aging population and the concomitant rise of aging-related diseases, a new interdisciplinary and practically orientated field like nutrgerontology is urgently needed.

## Funding

No funding information provided.

## Conflict of interest

The author has nothing to disclose.

## References

- Abuznait AH, Qosa H, Busnena BA, El Sayed KA, Kaddoumi A (2013) Olive-oil-derived oleocanthal enhances  $\beta$ -amyloid clearance as a potential neuroprotective mechanism against Alzheimer's disease: *in vitro* and *in vivo* studies. *ACS Chem. Neurosci.* **4**, 973–982.
- Alers S, Löffler AS, Wesselborg S, Stork B (2012) Role of AMPK-mTOR-Ulk1/2 in the regulation of autophagy: cross talk, shortcuts, and feedbacks. *Mol. Cell. Biol.* **32**, 2–11.
- Allen NE, Appleby PN, Davey GK, Kaaks R, Rinaldi S, Key TJ (2002) The associations of diet with serum insulin-like growth factor I and its main binding proteins in 292 women meat-eaters, vegetarians, and vegans. *Cancer Epidemiol. Biomarkers Prev.* **11**, 1441–1448.
- Ambrosone CB, McCann SE, Freudenheim JL, Marshall JR, Zhang Y, Shields PG (2004) Breast cancer risk in premenopausal women is inversely associated with consumption of broccoli, a source of isothiocyanates, but is not modified by GST genotype. *J. Nutr.* **134**, 1134–1138.
- Anon (2011) *Global Health and Aging*, World Health Organization. URL: [http://www.who.int/ageing/publications/global\\_health.pdf](http://www.who.int/ageing/publications/global_health.pdf). Last accessed on 2 June 2014
- Bartke A, Sun LY, Longo V (2013) Somatotrophic signaling: trade-offs between growth, reproductive development, and longevity. *Physiol. Rev.* **93**, 571–598.
- Barzilai N, Rennert G (2012) The rationale for delaying aging and the prevention of age-related diseases. *Rambam Maimonides Med. J.* **3**, e0020.
- Beulens JWW, de Bruijne LM, Stolk RP, Peeters PHM, Bots ML, Grobbee DE, van der Schouw YT (2007) High dietary glycemic load and glycemic index increase risk of cardiovascular disease among middle-aged women: a population-based follow-up study. *J. Am. Coll. Cardiol.* **50**, 14–21.
- Bjelakovic G, Nikolova D, Gluud LL, Simonetti RG, Gluud C (2007) Mortality in randomized trials of antioxidant supplements for primary and secondary prevention: systematic review and meta-analysis. *JAMA* **297**, 842–857.
- Blagosklonny MV (2006) Aging and immortality conceptual review. *Cell Cycle*, **5**, 2087–2102.

- Blüher M, Kahn BB, Kahn CR (2003) Extended longevity in mice lacking the insulin receptor in adipose tissue. *Science* **299**, 572–574.
- Boettler U, Sommerfeld K, Volz N, Pahlke G, Teller N, Somoza V, Lang R, Hofmann T, Marko D (2011) Coffee constituents as modulators of Nr2 nuclear translocation and ARE (EpRE)-dependent gene expression. *J. Nutr. Biochem.* **22**, 426–440.
- Bonafè M, Barbieri M, Marchegiani F, Olivieri F, Ragno E, Giampieri C, Mugianesi E, Centurelli M, Franceschi C, Paolisso G (2003) Polymorphic variants of insulin-like growth factor I (IGF-I) receptor and phosphoinositide 3-kinase genes affect IGF-I plasma levels and human longevity: cues for an evolutionarily conserved mechanism of life span control. *J. Clin. Endocrinol. Metab.* **88**, 3299–3304.
- Brouns F, Bjorck I, Frayn KN, Gibbs AL, Lang V, Slama G, Wolever TMS (2005) Glycaemic index methodology. *Nutr. Res. Rev.* **18**, 145–171.
- Bruning A (2013) Inhibition of mTOR signaling by quercetin in cancer treatment and prevention. *Anticancer Agents Med. Chem.* **13**, 1025–1031.
- Buitrago-Lopez A, Sanderson J, Johnson L, Warnakula S, Wood A, Di Angelantonio E, Franco OH (2011) Chocolate consumption and cardiometabolic disorders: systematic review and meta-analysis. *BMJ* **343**, d4488.
- Calabrese EJ, Bachmann KA, Bailer AJ, Bolger PM, Borak J, Cai L, Cedergreen N, Cherian MG, Chiueh CC, Clarkson TW, Cook RR, Diamond DM, Doolittle DJ, Dorato MA, Duke SO, Feinendegen L, Gardner DE, Hart RW, Hastings KL, Hayes AW, Hoffmann GR, Ives JA, Jaworowski Z, Johnson TE, Jonas WB, Kaminski NE, Keller JG, Klaunig JE, Knudsen TB, Kozumbo WJ, Lettieri T, Liu S-Z, Maisseu A, Maynard KI, Masoro EJ, McClellan RO, Mehendale HM, Mothersill C, Newlin DB, Nigg HN, Oehme FW, Phalen RF, Philbert MA, Rattan SJS, Riviere JE, Rodricks J, Sapolsky RM, Scott BR, Seymour C, Sinclair DA, Smith-Sonneborn J, Snow ET, Spear L, Stevenson DE, Thomas Y, Tubiana M, Williams GM, Mattson MP (2007) Biological stress response terminology: Integrating the concepts of adaptive response and preconditioning stress within a hormetic dose-response framework. *Toxicol. Appl. Pharmacol.* **222**, 122–128.
- Castaneda C, Gordon PL, Fielding RA, Evans WJ, Crim MC (2000) Marginal protein intake results in reduced plasma IGF-I levels and skeletal muscle fiber atrophy in elderly women. *J. Nutr. Health Aging* **4**, 85–90.
- Cavuto P, Fenech MF (2012) A review of methionine dependency and the role of methionine restriction in cancer growth control and life-span extension. *Cancer Treat. Rev.* **38**, 726–736.
- Cheng C-W, Adams GB, Perin L, Wei M, Zhou X, Lam BS, Da Sacco S, Mirisola M, Quinn DI, Dorff TB, Kopchick JJ, Longo VD (2014) Prolonged fasting reduces IGF-1/PKA to promote hematopoietic-stem-cell-based regeneration and reverse immunosuppression. *Cell Stem Cell* **14**, 810–823.
- Choi AMK, Ryter SW, Levine B (2013) Autophagy in human health and disease. *N. Engl. J. Med.* **368**, 651–662.
- Clayton RN (2003) Cardiovascular function in acromegaly. *Endocr. Rev.* **24**, 272–277.
- Colman RJ, Anderson RM, Johnson SC, Kastman EK, Kosmatka KJ, Beasley TM, Allison DB, Cruzen C, Simmons HA, Kemnitz JW, Weindruch R (2009) Caloric restriction delays disease onset and mortality in rhesus monkeys. *Science* **325**, 201–204.
- Colman RJ, Beasley TM, Kemnitz JW, Johnson SC, Weindruch R, Anderson RM (2014) Caloric restriction reduces age-related and all-cause mortality in rhesus monkeys. *Nat. Commun.* **5**, 3557.
- Cuervo AM (2008) Autophagy and aging: keeping that old broom working. *Trends Genet.* **24**, 604–612.
- De Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, Guidollet J, Touboul P, Delaye J (1994) Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* **343**, 1454–1459.
- Desideri G, Kwik-Urbe C, Grassi D, Necozione S, Ghiadoni L, Mastroiacovo D, Raffaele A, Ferri L, Bocale R, Lechiara MC, Marini C, Ferri C (2012) Benefits in cognitive function, blood pressure, and insulin resistance through cocoa flavanol consumption in elderly subjects with mild cognitive impairment: the Cocoa, Cognition, and Aging (CoCoA) study. *Hypertension* **60**, 794–801.
- Ebbeling CB, Swain JF, Feldman HA, Wong WW, Hachey DL, Garcia-Lago E, Ludwig DS (2012) Effects of dietary composition on energy expenditure during weight-loss maintenance. *JAMA* **307**, 2627–2634.
- Efeyan A, Zoncu R, Sabatini DM (2012) Amino acids and mTORC1: from lysosomes to disease. *Trends Mol. Med.* **18**, 524–533.
- Ericson U, Sonestedt E, Gullberg B, Hellstrand S, Hindy G, Wirfält E, Orho-Melander M (2013) High intakes of protein and processed meat associate with increased incidence of type 2 diabetes. *Br. J. Nutr.* **109**, 1143–1153.
- Fau D, Peret J, Hadjiiski P (1988) Effects of ingestion of high protein or excess methionine diets by rats for two years. *J. Nutr.* **118**, 128–133.
- Fontana L, Meyer TE, Klein S, Holloszy JO (2004) Long-term calorie restriction is highly effective in reducing the risk for atherosclerosis in humans. *Proc. Natl Acad. Sci. USA* **101**, 6659–6663.
- Fontana L, Partridge L, Longo VD (2010) Extending healthy life span—from yeast to humans. *Science* **328**, 321–326.
- Foster-Powell K, Holt SHA, Brand-Miller JC (2002) International table of glycemic index and glycemic load values: 2002. *Am. J. Clin. Nutr.* **76**, 5–56.
- Frydman-Marom A, Levin A, Farfara D, Benromano T, Scherzer-Attali R, Peled S, Vassar R, Segal D, Gazit E, Frenkel D, Ovadia M (2011) Orally administered cinnamon extract reduces  $\beta$ -amyloid oligomerization and corrects cognitive impairment in Alzheimer's disease animal models. *PLoS ONE* **6**, e16564.
- Gems D, de la Guardia Y (2012) Alternative perspectives on aging in *Caenorhabditis elegans*: reactive oxygen species or hyperfunction? *Antioxid. Redox Signal.* **00**, 1–9.
- Glantschnig H, Fisher JE, Wesolowski G, Rodan GA, Reszka AA (2003) M-CSF, TNF $\alpha$  and RANK ligand promote osteoclast survival by signaling through mTOR/S6 kinase. *Cell Death Differ.* **10**, 1165–1177.
- Gonzalez CA, Riboli E (2010) Diet and cancer prevention: contributions from the European Prospective Investigation into Cancer and Nutrition (EPIC) study. *Eur. J. Cancer* **46**, 2555–2562.
- Grandison RC, Piper MDW, Partridge L (2009) Amino-acid imbalance explains extension of lifespan by dietary restriction in *Drosophila*. *Nature* **462**, 1061–1064.
- Guevara-Aguirre J, Balasubramanian P, Guevara-Aguirre M, Wei M, Madia F, Cheng C-W, Hwang D, Martin-Montalvo A, Saavedra J, Ingles S, de Cabo R, Cohen P, Longo VD (2011) Growth hormone receptor deficiency is associated with a major reduction in pro-aging signaling, cancer, and diabetes in humans. *Sci. Transl. Med.* **3**, 70ra13.
- Halberg N, Henriksen M, Söderhamn N, Stallknecht B, Ploug T, Schjerling P, Dela F (2005) Effect of intermittent fasting and refeeding on insulin action in healthy men. *J. Appl. Physiol.* **99**, 2128–2136.
- Hansen M, Flatt T, Aguilaniu H (2013) Reproduction, fat metabolism, and life span: what is the connection? *Cell Metab.* **17**, 10–19.
- Harrison DE, Strong R, Sharp ZD, Nelson JF, Astle CM, Flurkey K, Nadon NL, Wilkinson JE, Frenkel K, Carter CS, Pahor M, Javors MA, Fernandez E, Miller RA (2009) Rapamycin fed late in life extends lifespan in genetically heterogeneous mice. *Nature* **460**, 392–395.
- Hayes JD, Kelleher MO, Eggleston IM (2008) The cancer chemopreventive actions of phytochemicals derived from glucosinolates. *Eur. J. Nutr.* **47**(Suppl 2), 73–88.
- Heilbronn LK, de Jonge L, Frisard MI, DeLany JP, Larson-Meyer DE, Rood J, Nguyen T, Martin CK, Volaufova J, Most MM, Greenway FL, Smith SR, Deutsch WA, Williamson DA, Ravussin E (2006) Effect of 6-month calorie restriction on biomarkers of longevity, metabolic adaptation, and oxidative stress in overweight individuals: a randomized controlled trial. *JAMA* **295**, 1539–1548.
- Holt SH, Miller JC, Petocz P (1997) An insulin index of foods: the insulin demand generated by 1000-kJ portions of common foods. *Am. J. Clin. Nutr.* **66**, 1264–1276.
- Holzenberger M, Dupont J, Ducos B, Leneuve P, Gélouën A, Even PC, Cervera P, Le Bouc Y (2003) IGF-1 receptor regulates lifespan and resistance to oxidative stress in mice. *Nature* **421**, 182–187.
- Hou NS, Taubert S (2012) Function and regulation of lipid biology in *Caenorhabditis elegans* aging. *Front. Physiol.* **3**, 143.
- Houston DK, Nicklas BJ, Ding J, Harris TB, Tyllavsky FA, Newman AB, Lee JS, Shayoun NR, Visser M, Kritchevsky SB (2008) Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) Study. *Am. J. Clin. Nutr.* **87**, 150–155.
- Houtkooper RH, Williams RW, Auwerx J (2010) Metabolic networks of longevity. *Cell* **142**, 9–14.
- Houtkooper RH, Pirinen E, Auwerx J (2012) Sirtuins as regulators of metabolism and healthspan. *Nat. Rev. Mol. Cell Biol.* **13**, 225–238.
- Hu R, Xu C, Shen G, Jain MR, Khor TO, Gopalkrishnan A, Lin W, Reddy B, Chan JY, Kong A-NT (2006) Gene expression profiles induced by cancer chemopreventive isothiocyanate sulforaphane in the liver of C57BL/6J mice and C57BL/6J/Nrf2 (-/-) mice. *Cancer Lett.* **243**, 170–192.
- Iwasaki K, Gleiser CA, Masoro EJ, McMahan CA, Seo EJ, Yu BP (1988) The influence of dietary protein source on longevity and age-related disease processes of Fischer rats. *J. Gerontol.* **43**, B5–B12.
- Jewell JL, Guan K-L (2013) Nutrient signaling to mTOR and cell growth. *Trends Biochem. Sci.* **38**, 233–242.
- Johnson SC, Rabinovitch PS, Kaerberlein M (2013) mTOR is a key modulator of ageing and age-related disease. *Nature* **493**, 338–345.
- Kanfi Y, Naiman S, Amir G, Peshti V, Zinman G, Nahum L, Bar-Joseph Z, Cohen HY (2012) The sirtuin SIRT6 regulates lifespan in male mice. *Nature* **483**, 218–221.
- Kapahi P, Zid BM, Harper T, Koslover D, Sapin V, Benzer S (2004) Regulation of lifespan in *Drosophila* by modulation of genes in the TOR signaling pathway. *Curr. Biol.* **14**, 885–890.
- Kenyon CJ (2010) The genetics of ageing. *Nature* **464**, 504–512.

- Kilhovd BK, Berg TJ, Birkeland KI, Thorsby P, Hanssen KF (1999) Serum levels of advanced glycation end products are increased in patients with type 2 diabetes and coronary heart disease. *Diabetes Care* **22**, 1543–1548.
- Kojima T, Kamei H, Aizu T, Arai Y, Takayama M, Nakazawa S, Ebihara Y, Inagaki H, Masui Y, Gondo Y, Sakaki Y, Hirose N (2004) Association analysis between longevity in the Japanese population and polymorphic variants of genes involved in insulin and insulin-like growth factor 1 signaling pathways. *Exp. Gerontol.* **39**, 1595–1598.
- Krebs M, Krssak M, Bernroider E, Anderwald C, Brehm A, Meyerspeer M, Nowotny P, Roth E, Waldhäusl W, Roden M (2002) Mechanism of amino acid-induced skeletal muscle insulin resistance in humans. *Diabetes* **51**, 599–605.
- Lee C, Raffaghello L, Brandhorst S, Safdie FM, Bianchi G, Martin-Montalvo A, Pistoia V, Wei M, Hwang S, Merlino A, Emionite L, de Cabo R, Longo VD (2012) Fasting cycles retard growth of tumors and sensitize a range of cancer cell types to chemotherapy. *Sci. Transl. Med.* **4**, 124ra27.
- Leto S, Kokkonen GC, Barrows CH (1976) Dietary protein, life-span, and biochemical variables in female mice. *J. Gerontol.* **31**, 144–148.
- Levine ME, Suarez JA, Brandhorst S, Balasubramanian P, Cheng C-W, Madia F, Fontana L, Mirisola MG, Guevara-Aguirre J, Wan J, Passarino G, Kennedy BK, Wei M, Cohen P, Crimmins EM, Longo VD (2014) Low protein intake is associated with a major reduction in IGF-1, cancer, and overall mortality in the 65 and younger but not older population. *Cell Metab.* **19**, 407–417.
- Lim EL, Hollingsworth KG, Aribisala BS, Chen MJ, Mathers JC, Taylor R (2011) Reversal of type 2 diabetes: normalisation of beta cell function in association with decreased pancreas and liver triacylglycerol. *Diabetologia* **54**, 2506–2514.
- Liu S, Willett WC, Stampfer MJ, Hu FB, Franz M, Sampson L, Hennekens CH, Manson JE (2000) A prospective study of dietary glycemic load, carbohydrate intake, and risk of coronary heart disease in US women. *Am. J. Clin. Nutr.* **71**, 1455–1461.
- Liu M, Wilk SA, Wang A, Zhou L, Wang R-H, Ogawa W, Deng C, Dong LQ, Liu F (2010) Resveratrol inhibits mTOR signaling by promoting the interaction between mTOR and DEPTOR. *J. Biol. Chem.* **285**, 36387–36394.
- López-Torres M, Barja G (2008) Lowered methionine ingestion as responsible for the decrease in rodent mitochondrial oxidative stress in protein and dietary restriction possible implications for humans. *Biochim. Biophys. Acta* **1780**, 1337–1347.
- Lu T, Sheng H, Wu J, Cheng Y, Zhu J, Chen Y (2012) Cinnamon extract improves fasting blood glucose and glycosylated hemoglobin level in Chinese patients with type 2 diabetes. *Nutr. Res.* **32**, 408–412.
- Martinet W, De Loof H, De Meyer GRY (2014) mTOR inhibition: a promising strategy for stabilization of atherosclerotic plaques. *Atherosclerosis* **233**, 601–607.
- Martini M, De Santis M, Braccini L, Gulluni F, Hirsch E (2014) PI3K/AKT signaling pathway and cancer: an updated review. *Ann. Med.* **46**, 1–12.
- Mattison JA, Roth GS, Beasley TM, Tilmont EM, Handy AM, Herbert RL, Longo DL, Allison DB, Young JE, Bryant M, Barnard D, Ward WF, Qi W, Ingram DK, de Cabo R (2012) Impact of caloric restriction on health and survival in rhesus monkeys from the NIA study. *Nature* **489**, 318–321.
- McCarty MF, Barroso-Aranda J, Contreras F (2009) The low-methionine content of vegan diets may make methionine restriction feasible as a life extension strategy. *Med. Hypotheses* **72**, 125–128.
- Meng X, Zhu K, Devine A, Kerr DA, Binns CW, Prince RL (2009) A 5-year cohort study of the effects of high protein intake on lean mass and BMC in elderly postmenopausal women. *J. Bone Miner. Res.* **24**, 1827–1834.
- Meyer KA, Kushi LH, Jacobs DR, Slavin J, Sellers TA, Folsom AR (2000) Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *Am. J. Clin. Nutr.* **71**, 921–930.
- Meyer TE, Kovács SJ, Ehsani AA, Klein S, Holloszy JO, Fontana L (2006) Long-term caloric restriction ameliorates the decline in diastolic function in humans. *J. Am. Coll. Cardiol.* **47**, 398–402.
- Mirisola MG, Taormina G, Fabrizio P, Wei M, Hu J, Longo VD (2014) Serine- and threonine/valine-dependent activation of PDK and Tor orthologs converge on Sch9 to promote aging. *PLoS Genet.* **10**, e1004113.
- Monti MC, Margarucci L, Tosco A, Riccio R, Casapullo A (2011) New insights on the interaction mechanism between tau protein and oleocanthal, an extra-virgin olive-oil bioactive component. *Food Funct.* **2**, 423–428.
- Mosdøl A, Witte DR, Frost G, Marmot MG, Brunner EJ (2007) Dietary glycemic index and glycemic load are associated with high-density-lipoprotein cholesterol at baseline but not with increased risk of diabetes in the Whitehall II study. *Am. J. Clin. Nutr.* **86**, 988–994.
- Muraki I, Imamura F, Manson JE, Hu FB, Willett WC, van Dam RM, Sun Q (2013) Fruit consumption and risk of type 2 diabetes: results from three prospective longitudinal cohort studies. *BMJ* **347**, f5001.
- O'Neill C (2013) PI3-kinase/Akt/mTOR signaling: impaired on/off switches in aging, cognitive decline and Alzheimer's disease. *Exp. Gerontol.* **48**, 647–653.
- O'Neill C, Kiely AP, Coakley MF, Manning S, Long-Smith CM (2012) Insulin and IGF-1 signalling: longevity, protein homeostasis and Alzheimer's disease. *Biochem. Soc. Trans.* **40**, 721–727.
- Ono K, Hasegawa K, Naiki H, Yamada M (2004) Curcumin has potent anti-amyloidogenic effects for Alzheimer's beta-amyloid fibrils *in vitro*. *J. Neurosci. Res.* **75**, 742–750.
- Orlich MJ, Singh PN, Sabaté J, Jaceldo-Siegl K, Fan J, Knutsen S, Beeson WL, Fraser GE (2013) Vegetarian dietary patterns and mortality in Adventist Health Study 2. *JAMA Intern. Med.* **173**, 1230–1238.
- Pan A, Sun Q, Bernstein AM, Schulze MB, Manson JE, Stampfer MJ, Willett WC, Hu FB (2012) Red meat consumption and mortality: results from 2 prospective cohort studies. *Arch. Intern. Med.* **172**, 555–563.
- Parrella E, Maxim T, Maialetti F, Zhang L, Wan J, Wei M, Cohen P, Fontana L, Longo VD (2013) Protein restriction cycles reduce IGF-1 and phosphorylated Tau, and improve behavioral performance in an Alzheimer's disease mouse model. *Aging Cell* **12**, 257–268.
- Partridge L (2012) Diet and healthy aging. *N. Engl. J. Med.* **367**, 2550–2551.
- Partridge L, Gems D (2007) Benchmarks for ageing studies. *Nature* **450**, 165–167.
- Pawlikowska L, Hu D, Huntsman S, Sung A, Chu C, Chen J, Joyner AH, Schork NJ, Hsueh W-C, Reiner AP, Psaty BM, Atzmon G, Barzilai N, Cummings SR, Browner WS, Kwok P-Y, Ziv E (2009) Association of common genetic variation in the insulin/IGF1 signaling pathway with human longevity. *Aging Cell* **8**, 460–472.
- Pedersen AN, Cederholm T (2014) Health effects of protein intake in healthy elderly populations: a systematic literature review. *Food Nutr. Res.* **58** <http://dx.doi.org/10.3402/fnr.v58.23364>.
- Price NL, Gomes AP, Ling AJY, Duarte FV, Martin-Montalvo A, North BJ, Agarwal B, Ye L, Ramadori G, Teodoro JS, Hubbard BP, Varela AT, Davis JG, Varamini B, Hafner A, Moaddel R, Rolo AP, Coppari R, Palmeira CM, de Cabo R, Baur JA, Sinclair DA (2012) SIRT1 is required for AMPK activation and the beneficial effects of resveratrol on mitochondrial function. *Cell Metab.* **15**, 675–690.
- Raffaghello L, Lee C, Safdie FM, Wei M, Madia F, Bianchi G, Longo VD (2008) Starvation-dependent differential stress resistance protects normal but not cancer cells against high-dose chemotherapy. *Proc. Natl Acad. Sci. USA* **105**, 8215–8220.
- Reinke A, Chen JC-Y, Aronova S, Powers T (2006) Caffeine targets TOR complex I and provides evidence for a regulatory link between the FRB and kinase domains of Tor1p. *J. Biol. Chem.* **281**, 31616–31626.
- Richie JP, Leutzinger Y, Parthasarathy S, Malloy V, Orentreich N, Zimmerman JA (1994) Methionine restriction increases blood glutathione and longevity in F344 rats. *FASEB J.* **8**, 1302–1307.
- Rohrmann S, Overvad K, Bueno-de-Mesquita HB, Jakobsen MU, Egeberg R, Tjønneland A, Nailler L, Boutron-Ruault M-C, Clavel-Chapelon F, Krogh V, Palli D, Panico S, Tumino R, Ricceri F, Bergmann MM, Boeing H, Li K, Kaaks R, Khaw K-T, Wareham NJ, Crowe FL, Key TJ, Naska A, Trichopoulou A, Trichopoulos D, Leenders M, Peeters PH, Engeset D, Parr CL, Skeie G, Jakyszyn P, Sanchez M-J, Huerta JM, Redondo ML, Barricarte A, Amiano P, Drake I, Sonestedt E, Hallmans G, Johansson I, Fedirko V, Romieux I, Ferrari P, Norat T, Vergnaud AC, Riboli E, Linseisen J (2013) Meat consumption and mortality - results from the European prospective investigation into cancer and nutrition. *BMC Med.* **11**, 63.
- Ross M (1961) Length of life and nutrition in the rat. *J. Nutr.* **75**, 197–210.
- Ross M, Bras G (1975) Food preference and length of life. *Science* **190**, 165–167.
- Salmerón J, Ascherio A, Rimm EB, Colditz GA, Spiegelman D, Jenkins DJ, Stampfer MJ, Wing AL, Willett WC (1997) Dietary fiber, glycemic load, and risk of NIDDM in men. *Diabetes Care* **20**, 545–550.
- Selman C, Lingard S, Choudhury AI, Batterham RL, Claret M, Clements M, Ramadan R, Okkenhaug K, Schuster E, Blanc E, Piper MD, Al-Qassab H, Speakman JR, Carmignac D, Robinson ICA, Thornton JM, Gems D, Partridge L, Withers DJ (2008) Evidence for lifespan extension and delayed age-related biomarkers in insulin receptor substrate 1 null mice. *FASEB J.* **22**, 807–818.
- Shah OJ, Wang Z, Hunter T (2004) Inappropriate activation of the TSC/Rheb/mTOR/S6K cassette induces IRS1/2 depletion, insulin resistance, and cell survival deficiencies. *Curr. Biol.* **14**, 1650–1656.
- Sieri S, Brighenti F, Agnoli C, Grioni S, Masala G, Bendinelli B, Sacerdote C, Ricceri F, Tumino R, Giurdanella MC, Pala V, Berrino F, Mattiello A, Chiodini P, Panico S, Krogh V (2013) Dietary glycemic load and glycemic index and risk of cerebrovascular disease in the EPICOR cohort. *PLoS ONE* **8**, e62625.
- Singh R, Barden A, Mori T, Beilin L (2001) Advanced glycation end-products: a review. *Diabetologia* **44**, 129–146.
- Solon-Biet SM, McMahon AC, Ballard JWO, Ruohonen K, Wu LE, Cogger VC, Warren A, Huang X, Pichaud N, Melvin RG, Gokarn R, Khalil M, Turner N, Cooney GJ, Sinclair DA, Raubenheimer D, Le Couteur DG, Simpson SJ (2014)

- The ratio of macronutrients, not caloric intake, dictates cardiometabolic health, aging, and longevity in *ad libitum*-fed mice. *Cell Metab.* **19**, 418–430.
- Steuerman R, Shevah O, Laron Z (2011) Congenital IGF1 deficiency tends to confer protection against post-natal development of malignancies. *Eur. J. Endocrinol.* **164**, 485–489.
- Stout MB, Tchkonja T, Pirtskhalava T, Palmer AK, List EO, Berryman DE, Lubbers ER, Escande C, Spong A, Masternak MM, Oberg AL, LeBrasseur NK, Miller RA, Kopchick JJ, Bartke A, Kirkland JL (2014) Growth hormone action predicts age-related white adipose tissue dysfunction and senescent cell burden in mice. *Aging* **6**, 575–586.
- Suh Y, Atzmon G, Cho M-O, Hwang D, Liu B, Leahy DJ, Barzilai N, Cohen P (2008) Functionally significant insulin-like growth factor I receptor mutations in centenarians. *Proc. Natl Acad. Sci. USA* **105**, 3438–3442.
- Taguchi A, Wartschow LM, White MF (2007) Brain IRS2 signaling coordinates life span and nutrient homeostasis. *Science* **317**, 369–372.
- Tchkonja T, Morbeck DE, Von Zglinicki T, Van Deursen J, Lustgarten J, Scrbale H, Khosla S, Jensen MD, Kirkland JL (2010) Fat tissue, aging, and cellular senescence. *Aging Cell* **9**, 667–684.
- Thomas DE, Elliott EJ, Baur L (2007) Low glycaemic index or low glycaemic load diets for overweight and obesity. *Cochrane Database Syst. Rev.*, CD005105.
- Tremblay F, Krebs M, Dombrowski L, Brehm A, Bernroider E, Roth E, Nowotny P, Waldhäusl W, Marette A, Roden M (2005) Overactivation of S6 kinase 1 as a cause of human insulin resistance during increased amino acid availability. *Diabetes* **54**, 2674–2684.
- Tremblay F, Brûlé S, Hee Um S, Li Y, Masuda K, Roden M, Sun XJ, Krebs M, Polakiewicz RD, Thomas G, Marette A (2007) Identification of IRS-1 Ser-1101 as a target of S6K1 in nutrient- and obesity-induced insulin resistance. *Proc. Natl Acad. Sci. USA* **104**, 14056–14061.
- Trinh K, Moore K, Wes PD, Muchowski PJ, Dey J, Andrews L, Pallanck LJ (2008) Induction of the phase II detoxification pathway suppresses neuron loss in *Drosophila* models of Parkinson's disease. *J. Neurosci.* **28**, 465–472.
- Um SH, D'Alessio D, Thomas G (2006) Nutrient overload, insulin resistance, and ribosomal protein S6 kinase 1, S6K1. *Cell Metab.* **3**, 393–402.
- Van Aller GS, Carson JD, Tang W, Peng H, Zhao L, Copeland RA, Tummino PJ, Luo L (2011) Epigallocatechin gallate (EGCG), a major component of green tea, is a dual phosphoinositide-3-kinase/mTOR inhibitor. *Biochem. Biophys. Res. Commun.* **406**, 194–199.
- Varady KA, Bhutani S, Church EC, Klempel MC (2009) Short-term modified alternate-day fasting: a novel dietary strategy for weight loss and cardioprotection in obese adults. *Am. J. Clin. Nutr.* **90**, 1138–1143.
- Vellai T, Takacs-Vellai K, Zhang Y, Kovacs AL, Orosz L, Müller F (2003) Genetics: influence of TOR kinase on lifespan in *C. elegans*. *Nature* **426**, 620.
- Villegas R, Liu S, Gao Y-T, Yang G, Li H, Zheng W, Shu XO (2007) Prospective study of dietary carbohydrates, glycemic index, glycemic load, and incidence of type 2 diabetes mellitus in middle-aged Chinese women. *Arch. Intern. Med.* **167**, 2310–2316.
- Vitek MP, Bhattacharya K, Glendening JM, Stopa E, Vlassara H, Bucala R, Manogue K, Cerami A (1994) Advanced glycation end products contribute to amyloidosis in Alzheimer disease. *Proc. Natl Acad. Sci. USA* **91**, 4766–4770.
- Wanke V, Cameroni E, Uotila A, Piccolis M, Urban J, Loewith R, De Virgilio C (2008) Caffeine extends yeast lifespan by targeting TORC1. *Mol. Microbiol.* **69**, 277–285.
- Willcox DC, Willcox BJ, Todoriki H, Curb JD, Suzuki M (2006) Caloric restriction and human longevity: what can we learn from the Okinawans? *Biogerontology* **7**, 173–177.
- Willcox BJ, Willcox DC, Todoriki H, Fujiyoshi A, Yano K, He Q, Curb JD, Suzuki M (2007) Caloric restriction, the traditional Okinawan diet, and healthy aging: the diet of the world's longest-lived people and its potential impact on morbidity and life span. *Ann. N. Y. Acad. Sci.* **1114**, 434–455.
- Willis MS, Patterson C (2013) Proteotoxicity and cardiac dysfunction—Alzheimer's disease of the heart? *N. Engl. J. Med.* **368**, 455–464.
- Winder WW, Hardie DG (1999) AMP-activated protein kinase, a metabolic master switch: possible roles in type 2 diabetes. *Am. J. Physiol.* **277**, E1–E10.
- Wu JJ, Liu J, Chen EB, Wang JJ, Cao L, Narayan N, Fergusson MM, Rovira II, Allen M, Springer DA, Lago CU, Zhang S, DuBois W, Ward T, de Cabo R, Gavrilova O, Mock B, Finkel T (2013) Increased mammalian lifespan and a segmental and tissue-specific slowing of aging after genetic reduction of mTOR expression. *Cell Rep.* **4**, 913–920.
- Wullschlegel S, Loewith R, Hall MN (2006) TOR signaling in growth and metabolism. *Cell* **124**, 471–484.
- Yang F, Lim GP, Begum AN, Ubada OJ, Simmons MR, Ambegaokar SS, Chen PP, Kaye R, Glabe CG, Frautschy SA, Cole GM (2005) Curcumin inhibits formation of amyloid beta oligomers and fibrils, binds plaques, and reduces amyloid *in vivo*. *J. Biol. Chem.* **280**, 5892–5901.
- Yu BP, Masoro EJ, McMahan CA (1985) Nutritional influences on aging of Fischer 344 rats: I. Physical, metabolic, and longevity characteristics. *J. Gerontol.* **40**, 657–670.